arXiv:cond-mat/9908422v1 28 Aug 1999

On the problem of an electron scattering in an arbitrary one-dimensional potential field

D.M. Sedrakian*, A.Zh.Khachatrian**

*Department of Physics, Yerevan State University,
Manoogian 1,Yerevan 375049, Armenia

**Department of General Physics, Armenian State Engineering
University, Terian 3, Yerevan 375046, Armenia

February 1, 2008

Abstract

Recurrent representations for an electron transmission and reflection amplitudes for a one-dimensional chain are obtained. The linear differential equations for scattering amplitudes of an arbitrary potential are found.

In this work the electron scattering problem is considered, when the potential energy has the form [1, 2, 3, 4, 5].

$$V(x) = \sum_{n=1}^{N} V_n(x - x_n) \tag{1}$$

Here $V(x - x_n)$ is an individual potential, which has located near the $x_n$point. It is suggested, that individual potentials $V(x - x_n)$ do not have general points.

There have been many studies to investigate electronic properties of one-dimensional quaziperiodical and disordered systems. It is well known, that the problem of determination of transmission amplitude (TA) and reflection amplitude (RA) in the field of (1) type is reduced to the problem of N matrixes product calculation [6, 7, 8];

$$\begin{pmatrix} T_N \\ 0 \end{pmatrix} = \prod_{N}^{n=1} \begin{pmatrix} 1/t_n^* & -r_n^*/t_n^* \\ -r_n/t_n & 1/t_n \end{pmatrix} \begin{pmatrix} 1 \\ R_N \end{pmatrix} \tag{2}$$

where T_N, $R_N$are TA and RA of potential $V(x)$, t_n and $r_n$are TA and RA of the individual potential $V(x-x_n)$ of the system (1).

The problem of calculation of the N two-order matrixes product (2), in general, is equivalent to the problem of solution of some linear equations [9]. So, the problem is represented as

$$D_N = r_N/t_N P_{N-1} + 1/t_N D_{N-1} N \geq 1 \tag{3}$$

$$P_N = r_N^*/t_N^* D_{N-1} + 1/t_N^* P_{N-1}.N \geq 1 \tag{4}$$

Here $D_N = 1/T_N$, $P_N = R_N^*/T_N^*$ and $D_{N-n} = 1/T_{N-n}$, $P_{N-n} = R_{N-n}^*/T_{N-n}^*$, where T_{N-n} and $R_{N-n}$are TA and RA of the first potentials of the field (1). Note, that in (3, 4) N is a variable.

Let us now introduce (3, 4) in more confortable form. Excluding P_{N-1} from the first equation and D_{N-1} from the second one, we will get two equations for D_N and P_N ;

$$D_N = A_N D_{N-1} - B_N D_{N-2} N \geq 2 \tag{5}$$

$$P_N = A_N^* P_{N-1} - B_N^* P_{N-2}, \quad N \geq 2 \tag{6}$$

where $A_N = 1/t_N + B_N/t_{N-1}^*$ and $B_N = r_N t_{N-1}/r_{N-1} t_N$.

So the problem of determination of $T_N$and R_N is reduced to the problem of solution of the equations (5, 6). In order to have the simple solution of (5, 6), it is necessary to give initial conditions for equations (5) and (6). They are

$$D_1 = t_1^{-1}, D_0 = 1, P_1 = r_1^*/t_1^*, P_0 = 0. \tag{7}$$

Note, that representations (5, 6) take place for arbitrary potentials of the type (1). So, if we take $t_N = 1 + i\alpha_N/2k$ and $r_N = -i(\alpha_N/2k)\exp 2ikx_N$, then (5) will coincide with the result of [10], received by the "determinant" method for δ- potentials. In a case of layered system from homogeneous mediums, (5, 6) yields to the result of [11].

It is interesting to apply the obtained result (5, 6) to a simple case, when potentials $V(x-x_n)$ are located periodically and have the same form.

Then $t_n$and r_n $(n = 1, 2, ..., N)$ can be represented as $t_n = t_1$ and $r_n = r_1 \exp 2ik(x_n - x_1)$. In this case, according to (5, 6), it is clear that coefficients A_N and B$_N$become independent from N and the equations (5, 6) are easily solved [12].

Taking into account conditions (7) and denoting the systems period by a, for D_N and P_N the following expressions are obtained

$$D_N = \exp iNka \, \left\{ \cos N\beta + iIm(t_1^{-1} \exp(-ika)) \sin N\beta / \sin\beta \right\}, \qquad (8)$$

$$P_N = r_1^* / t_1^* \exp -i(N-1)ka \, \sin N\beta / \sin\beta \, .$$

Here $k^2 = E$ $(\hbar = 2m_0 = 1)$ is the electron energy. The electron energy spectrum is given by $\cos\beta = Re(t_1^{-1} \exp(-ika))$, which was obtained in [13].

Let us now considered one important form of the potential $V(x)$ (1), when $V(x - x_n)$ the are rectangular potentials with arbitrary width $2d_n$ and magnitudes of potentials V_n. Then $t_n$and r_n are given by well-known formulas [14];

$$t_n^{-1} = \exp i2kd_n \, \left\{ \cos 2kd_n - i\frac{k_n^2 + k^2}{2k_n k} \sin 2kd_n \right\}, \qquad (9)$$

$$r_n / t_n = i \exp i2kx_n \, \frac{k_n^2 - k^2}{2k_n k} \sin 2kd_n. \qquad (10)$$

where $k_n = \sqrt{E - V_n}$, x_n is a coordinate of the middle point of the potential.

This model (9, 10) can be used to obtain differential equations for $D(x) = 1/T(x)$ and $P(x) = R^*(x)/T^*(x)$ quantities, where $T(x)$ and $R(x)$ are TA and RA of the potential

$$V(x) = U(y)\theta(x - y) \qquad (11)$$

where $U(y)$ is an arbitrary function limited within some interval $a \leq y \leq b$ and is equal to zero outside of that.

Indeed, approximating $V(x)$ by rectangular potentials, it is possible with the help of representation (5, 6) and corresponding limit transition to get the sought equation. If we denote in (5, 6) $D_{N-1} = D(x)$, then $D_N = D(x + \Delta x)$ will correspond to the potential $V(x + \Delta x)$. Respectively, $D_{N-2} = D(x - \Delta x)$

will correspond to the potential $V(x-\Delta x)$. Presenting in the expression (9, 10,) $V_N = V(x+\Delta x), V_{N-1} = V(x), 2d_n = \Delta x$ and expanding (5, 6) in a series by infinitely small quantity Δx, we get

$$\frac{1+B}{2}\frac{d^2D}{dx^2}(\Delta x)^2 + (1-B)\frac{dD}{dx}\Delta x - (A-B-1)D + \theta\left((\Delta x)^3\right) = 0, \quad (12)$$

where

$$B-1 = \left(2ik + \frac{1}{V(x)}\frac{dV(x)}{dx}\right) * \Delta x + \theta\left((\Delta x)^2\right),$$

$$A-B-1 = V(x)(\Delta x)^2 + \theta\left((\Delta x)^3\right). \quad (13)$$

Inserting (13) in to (12) and taking $\Delta x \to 0$, we find the desired equation for $D(x)$:

$$\frac{d^2D}{dx^2} - \left(2ik + \frac{1}{V(x)}\frac{dV(x)}{dx}\right)\frac{dD}{dx} - V(x)D = 0 \quad (14)$$

To find TA and RA of the potential $V(x)$ it is necessary to solve the equation (14) with the following initial conditions:

$$D(a) = 1, dD/dx\mid_{x=a} = iV(a)/2k \quad (15)$$

The value of the function $D(x)$ in the point x determines TA of the potential $V(x)$. According to the equations (5, 6), the quantity $P^*(x)$ satisfies the obtained equation (14) too. The initial conditions for $P^*(x)$ are the following

$$P^*(a) = 0, dP^*/dx\mid_{x=a} = iV(a)/2k \exp i2ka \quad (16)$$

Let us now consider the obtained result (14) for a case of a simple rectangular potential wall with width $L = b-a$ and a constant potential $V(x) = V_n$. Then, it is easy to see, that the solution of the problem (14)-(15) is

$$D(b) = \exp ikL\ \left\{\cos k_nL - i\frac{k_n^2+k^2}{2k_nk}\sin k_nL\right\}, \quad (17)$$

which, as was expected, agrees with (9) for $L = 2d_n$.

In conclusion, it is interesting to note, that taking $\exp -ikxD(x)$ as $F_1(x)$ and $\exp -ikxP(x)$ as $F_2(x)$, from equation (14), we obtain the Schrödinger equation for function $F_1(x) - F_2(x) = L(x)$;

$$\frac{d^2L}{dx^2} + (E - V(x))\,L = 0 \tag{18}$$

while the $F_1(x) + F_2(x) = \frac{i}{k}\frac{dL(x)}{dx}$.

Thus, the problem of determination of TA and RA is reduced to the Cauchy problem for Shrodinger equation (18), with initial condition (7).

References

[1] I.M. Lifshits, S.A. Gredeskul, L.A. Pastur, Introduction to the Theory of Disordered System (Wiley, Now York, 1988).

[2] T. Hirota and K. Ishii, Prog. Theor. Phys. 45 (1971) 1713.

[3] D.J. Thouless, J.Phys. C5 (1972) 77.

[4] P.W. Anderson, D.J. Thouless, E. Abrahams, D.S. Fisher, Phys. Rev. B22 (1980) 3519.

[5] V.N. Prigodin, Sov. Phys. JETP 52 (1980) 1185.

[6] A. Abrikosov , Solid State Commun. 37 (1981) 997.

[7] P. Erdos, R.C. Herndon, Adv. Phys. 31 (1982) 65, Solid State Commun. 98 (1996) 495.

[8] M. Ya Azbel, Phys. Rev. B22 (1983) 4106 .

[9] D.M. Sedrakian, A. Zh. Khachatrian, Prog. NAS Armenia 34 (1999) 138.

[10] V.M. Gasparian, B.L. Altshuler, A.G. Aronov, Z.H. Kasamanian, Phys. Lett. A132 (1988) 201.

[11] A.G. Aronov, V.M. Gasparian, J Phys.: Condens.Matter 3 (1991) 3023.

[12] D.M. Sedrakian, D.H. Badalian, V.M. Gasparian, A. Zh. Khachatrian, Rus. Phys. JETP 82 (1996) 129.

[13] V. Heine, Group Theory in Quantum Mechanics (London-Paris, Pergamon Press, 1960).

[14] L.D. Landau and E.M. Lifshits, Quantum Mechanics (Moscow, 1974).